

AD _____

GRANT NO: DAMD17-94-J-4344

TITLE: Vitamin D and Breast Cancer

PRINCIPAL INVESTIGATOR: Esther Janowsky, M.D., M.P.H.

CONTRACTING ORGANIZATION: University of North Carolina
at Chapel Hill
Chapel Hill, NC 27599-4100

REPORT DATE: April 1997

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick
Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19970711 089

DTIC QUALITY INSPECTED 3

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

| | | | | |
|---|--|---|---|---------------------------|
| 1. AGENCY USE ONLY (Leave blank) | | 2. REPORT DATE April 1997 | 3. REPORT TYPE AND DATES COVERED Annual 1 Jul 95 - 30 Jun 96 | |
| 4. TITLE AND SUBTITLE Vitamin D and Breast Cancer | | | 5. FUNDING NUMBERS DAMD17-94-J-4344 | |
| 6. AUTHOR(S) Esther Janowsky, M.D., M.P.H. | | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of North Carolina at Chapel Hill Chapel Hill, North Carolina 27599-4100 | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 | | | 10. SPONSORING / MONITORING AGENCY REPORT NUMBER | |
| 11. SUPPLEMENTARY NOTES | | | | |
| 12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release, distribution unlimited | | | 12b. DISTRIBUTION CODE | |
| 13. ABSTRACT (Maximum 200 words) We analyzed 511 (case=156, control 1=184, control 2=171) archived samples of blood from a previous case-control study of breast cancer for two vitamin D metabolites: 25(OH)D and 1,25(OH) ₂ D. Our data indicate that white women at the time of first diagnosis of breast cancer have significantly lower mean levels of 1,25(OH) ₂ D than comparable women without breast cancer. The odds ratio (95% CI) for lowest relative to highest quartile of 1,25(OH) ₂ D was 6.3 (2.6, 15.3) in a comparison of white cases and control 1 subjects, after adjustment for age, month of blood draw, clinic, assay batch, and sample storage time. The risk was stronger in women above the median age of 54, 5.2 (2.4, 11.1), than for younger women, 1.3 (0.7, 2.6). We did not see the same pattern of risk associated with low 1,25(OH) ₂ D among the small number of black women in our study. There were no differences between cases and controls in mean 25(OH)D levels in either race. This work may have important implications for the chemoprevention and treatment of breast cancer. | | | | |
| 14. SUBJECT TERMS 1. Epidemiology 2. 25-hydroxyvitamin D 3. vitamin D receptors | | 4. 1,25-dihydroxyvitamin D 5. Vitamin D metabolites 6. Immunohistochemical assays 7. Breast Cancer | | 15. NUMBER OF PAGES 14 |
| 17. SECURITY CLASSIFICATION OF REPORT Unclassified | | 18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified | | 16. PRICE CODE |
| 19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified | | 20. LIMITATION OF ABSTRACT Unlimited | | |

FOREWORD


Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.


In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

 For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

 April 15 '87
PI - Signature Date

DTIC QUALITY INSPECTED 3

Table of Contents

Page

| | |
|----|------------------------------------|
| 1 | Front Cover |
| 2 | SF 298 (Report Documentation Page) |
| 3 | Foreword |
| 4 | Table of Contents |
| 5 | Introduction |
| 7 | Body |
| 12 | Conclusions |
| 13 | References |

INTRODUCTION

Recent work has defined a previously unsuspected involvement of vitamin D in cellular growth and differentiation^{1,2}. This recognition has fostered an interest in the investigation of a possible role for vitamin D in carcinogenesis³. Several cancers have been evaluated for a relationship between tumor occurrence and low levels of vitamin D. There is a suggestion in the literature that decreased intake of vitamin D may be associated with cancer of the colon⁴; blood levels of vitamin D are lower in African-American and white men with cancer of the prostate than in similar men without prostate cancer⁵.

Evidence for the possible relationship between vitamin D and breast cancer is based on several lines of investigation. Ecologic studies generally support a relationship between low levels of sunlight exposure and breast cancer^{6,7,8}. A single case-control study in Canada, however, failed to demonstrate an association between low levels of vitamin D, as determined by dietary history, and breast cancer⁹. Studies in rats show an increased incidence of mammary tumors under conditions of low dietary vitamin D (0.05 IU/kcal) and calcium (0.25 mg/kcal)¹⁰. 1,25-dihydroxy vitamin D (1,25(OH)₂D), which is the hormonally active metabolite, displays a growth-inhibitory effect on human breast adenocarcinoma cells in culture irrespective of their sex-steroid dependence¹¹. Cultures of the human breast cancer cell line BT-20 demonstrate increased differentiation when exposed to 1,25(OH)₂D daily for 8 days¹².

The purpose of our current work is to determine whether there are differences in blood levels of 1,25-dihydroxyvitamin D between women with breast cancer and two control groups of women without breast cancer. We are using archived samples of blood obtained from women in a just-completed case-control study of genetic risk factors for breast cancer. In addition, we are working to modify the immunohistochemical technique for detection of vitamin D receptors (VDR) described for frozen tissue for use in paraffin embedded tissue. If we are successful, we will be able to evaluate the VDR status of tumor tissue from our subjects with breast cancer.

The specific aims of our study are as follows: 1) To determine if blood levels of 25-hydroxy vitamin D and 1,25-dihydroxyvitamin D are lower in women at the time of first diagnosis of breast cancer than in comparable women who do not have breast cancer. 2) To describe the relationship between levels of 25-hydroxy vitamin D and 1,25-dihydroxyvitamin D and previously described risk factors for breast cancer. 3) To identify and quantify the distribution of vitamin D receptors in tumor samples from the women diagnosed with breast cancer. 4) To identify the relationship between previously described risk factors for breast cancer and blood levels of 25-hydroxy vitamin D and 1,25-dihydroxyvitamin D with vitamin D receptor (+) and vitamin D receptor (-) breast cancers.

Body

Task 1: Analysis of vitamin D metabolites (0-15) months.

We have completed the analysis of the vitamin D metabolites for the entire group of 511 samples. Our vitamin D assays were done in 13 batches of 40 samples each with an admixture of randomly selected samples consisting of approximately one-third cases and one-third controls from each of the two control groups. Fourteen percent of the samples had to be reassayed because of technical difficulties such as problems with recovery. We used the calf thymus radioreceptor assay from INCSTAR (Stillwater, Minnesota) for determination of 1,25(OH)₂D and the radioimmunoassay from INCSTAR (Stillwater, Minnesota) for determination of 25(OH)D. A summary of the mean levels of the precursor metabolite, 25(OH)D, and the active metabolite, 1,25(OH)₂D as well as median values and ranges by race are presented in Table 1. 8 Asian-American, 1 Native American, and 5 Hispanic women were excluded from the analyses because their numbers were too small to provide meaningful results.

Table 1. Comparison of vitamin D metabolite levels in cases and controls by race

| | case n=21 | Black control 1 n=30 | control 2 n=23 | case n=131 | White control 1 n=149 | control 2 n=143 |
|--------------|--------------|----------------------------|-------------------|---------------|-----------------------------|--------------------|
| D1,25, pg/ml | | | | | | |
| * mean (SD) | 23.87 (9.42) | 21.42 (8.68) | 25.73 (10.45) | 18.59 (8.62) | 23.53 (9.01) | 22.39 (9.02) |
| median | 21.47 | 20.10 | 26.80 | 17.18 | 22.29 | 20.63 |
| range | 6.71 - 45.16 | 5.28 - 39.49 | 6.35 - 44.07 | 2.71 - 45.29 | 1.88 - 47.51 | 4.61 - 50.85 |
| D25, ng/ml | | | | | | |
| * mean (SD) | 8.91 (6.50) | 9.94 (7.40) | 11.06 (6.42) | 15.06 (8.81) | 14.71 (7.41) | 13.96 (6.45) |
| median | 6.80 | 8.125 | 7.99 | 13.49 | 12.98 | 13.31 |
| range | 3.52 - 28.47 | 1.87 - 40.33 | 3.81 - 26.73 | 0 - 48.44 | 2.14 - 48.40 | 3.0 - 40.71 |

*P-value for t test of difference in means:

| | Black | White |
|-------|--|--|
| D1,25 | case versus control 1 0.34 case versus control 2 0.54 | case versus control 1 0.0001 case versus control 2 0.0004 |
| D25 | case versus control 1 0.61 case versus control 2 0.28 | case versus control 1 0.72 case versus control 2 0.24 |

There was a significant difference in 1,25(OH)₂D pg/ml between white cases and each control group; there were no differences in mean values of 25(OH)D. There were no

statistically significant differences in either 1,25(OH)₂D or 25(OH)D between black cases and either control group.

The assay of the vitamin D metabolites has been completed; in white women, the results support the hypothesis that women with breast cancer have lower levels of 1,25(OH)₂D at the time of first diagnosis of their disease than comparable women without breast cancer. We do not see the same relationship between 1,25(OH)₂D level and disease risk in the small number of black women in the study.

Task 2: Development of Assay for VDR (0-6 months)

We have made slow progress in developing the assay; we are using human intestine, known to have the VDR, for the development of the assay in our lab. We have been collaborating with Dr. Lynn Dressler in the Molecular Epidemiology Laboratory at the University of North Carolina. At present, we are using streptavidin conjugated to glucose oxidase. This has resulted in a marked decrease in background staining and better color development. After establishing reliable positive and negative controls, we will begin to use the assay in breast tissue.

In preparation for the application of these methods to fixed tissues, we have established a working relationship with Dr. Lester Layfield's laboratory (Department of Pathology, Duke University). In this clinical lab, they have adapted methods for the localization of the estrogen receptor in frozen tissues for use on fixed specimens. Dr. Layfield has offered several protocols for our use and the resources of his laboratory to enable us to successfully adapt the immunohistochemical methods for VDR visualization in frozen specimens for sections of fixed tissue.

Task 3: Detection of VDR's in paraffin-embedded tumor specimens (6-18 months).

We have been unable to proceed with this task because of the difficulties in developing a satisfactory assay for the VDR (see Task 2).

Task 4: Data analysis and report writing (15-24 months).

We have completed analysis of the data relating to Specific Aim 1 of our study: to determine if blood levels of 25(OH)D and 1,25(OH)₂D are lower in women at the time of initial diagnosis of breast cancer than in comparable women who do not have breast cancer. Table 2 describes the risk associated with being in various quartiles of 1,25(OH)₂D with the highest quartile as the reference. Black subjects did not show the pattern of increased risk associated with low levels of 1,25(OH)₂D that was evident in white subjects. Risk levels were highest for the comparison of white cases and control 1 subjects, groups who were matched on month of blood drawing. When adjusted for age, month of blood draw, clinic, assay batch, and sample storage time, the OR (95% CI) for lowest versus highest quartile of 1,25(OH)₂D for white cases and control 1 women was 6.3 (2.6, 15.3)

Table 2. Crude Odds ratios (95% CI) for the risk of breast cancer associated with quartile of 1,25(OH)₂D

| | n | Quartile of 1,25(OH) ₂ D pg/ml* | | | 4 (ref) | p trend |
|-----------------|-----|--|----------------|----------------|---------|----------|
| | | 1 | 2 | 3 | | |
| All subjects | 511 | 2.9 (1.7, 5.1) | 1.5 (0.8, 2.7) | 1.4 (0.8, 2.6) | 1 | < 0.0001 |
| Black subjects | 74 | 0.8 (0.2, 3.6) | 1.6 (0.4, 6.5) | 1.1 (0.3, 4.6) | 1 | 0.96 |
| White subjects | 423 | 3.8 (2.0, 7.0) | 1.5 (0.8, 3.1) | 1.4 (0.7, 2.8) | 1 | <0.0001 |
| cases/control 1 | 280 | 4.4 (2.2, 8.7) | 1.3 (0.6, 2.8) | 1.3 (0.6, 2.8) | 1 | <0.0001 |
| cases/control 2 | 274 | 3.4 (1.7, 6.9) | 1.6 (0.8, 3.4) | 1.8 (0.9, 3.9) | 1 | 0.001 |
| quartiles*: | | 1 | 2 | 3 | 4 | |
| all | | ≤16.470 | 16.48 - 21.38 | 21.39 - 28.04 | > 28.04 | |
| black | | ≤16.699 | 16.70 - 22.07 | 22.08 - 30.44 | >30.44 | |
| white | | ≤16.597 | 16.60 - 21.62 | 21.63 - 28.07 | >28.07 | |
| control 1 | | ≤18.130 | 18.14 - 22.29 | 22.29 - 27.94 | >27.94 | |
| control 2 | | ≤15.720 | 15.73 - 20.64 | 20.65 - 28.39 | >28.39 | |

*quartile ranges in picograms/ml are based on the distribution of metabolite in the respective control group; values for control groups 1 and 2 are based on white subjects

We have also completed the analysis of the data relating to Specific Aim 2: to describe the relationship between levels of 25(OH)D and 1,25(OH)₂D and previously described risk factors for breast cancer. There were no significant associations between either lower or higher levels of 25(OH)D and previously described risk factors for breast cancer such as older age, postmenopausal status, higher BMI, younger age at menarche, or use of hormone replacement therapy among the white control 1 women. 1,25(OH)₂D levels were higher in postmenopausal white control 1 women (p=0.046).

We did a stratified analysis with white subjects, cases and control group 1 to investigate possible interactions between 1,25(OH)₂D and selected covariates (Table 3).

Table 3. Stratified analysis of breast cancer risk associated with levels of 1,25(OH)₂D below the median value of 22.29 pg/ml, white cases and control group 1 (n=280)

| Variable | 1,25(OH) ₂ D pg/ml | | | | OR (95% CI) | p* |
|-----------------------------|-------------------------------|------|--------|------|-----------------|--------|
| | ≤ 22.29 | | >22.29 | | | |
| | case | cont | case | cont | | |
| Age, years | | | | | | |
| > 54 | 50 | 31 | 13 | 42 | 5.2 (2.4, 11.1) | 0.0008 |
| ≤ 54 | 43 | 43 | 25 | 33 | 1.3 (0.7, 2.6) | |
| Menopausal status | | | | | | |
| post- | 64 | 45 | 25 | 57 | 3.2 (1.8, 5.9) | 0.11 |
| pre- | 29 | 29 | 13 | 18 | 1.4 (0.6, 3.3) | |
| BMI, kg/m ² | | | | | | |
| > 24.13 | 44 | 32 | 21 | 42 | 2.8 (1.4, 5.5) | 0.70 |
| ≤ 24.13 | 49 | 42 | 17 | 33 | 2.3 (1.1, 4.6) | |
| 25(OH)D, ng/ml | | | | | | |
| ≤ 12.98 | 45 | 35 | 19 | 40 | 2.7 (1.3, 5.5) | 0.78 |
| > 12.98 | 48 | 39 | 19 | 35 | 2.3 (1.1, 4.6) | |
| Hormone replacement therapy | | | | | | |
| yes | 37 | 29 | 16 | 29 | 2.3 (1.1, 5.0) | 0.85 |
| no | 56 | 45 | 22 | 45 | 2.5 (1.3, 4.8) | |

*p value for the Breslow Day χ^2_{1df} test for homogeneity of the stratum specific ORs

An interaction was considered present if the Breslow-Day statistic for homogeneity of the odds ratio was significant at the 0.05 level. The magnitude of breast cancer risk related to low $1,25(\text{OH})_2\text{D}$ level was greater for women above the age of 54 than for women 54 and younger ($p=0.0008$). There was no difference in risk according to menopausal status, BMI, $25(\text{OH})\text{D}$ level, or use of hormone replacement therapy.

We have prepared and submitted a manuscript on the results of our study; we are preparing a second manuscript describing the methods used to assess assay validity and reliability for $25(\text{OH})\text{D}$ and $1,25(\text{OH})_2\text{D}$ in whole blood. Preparation of the Final Report for the DOD is underway.

Conclusion

Our data indicate that white women in the lowest quartile of $1,25(\text{OH})_2\text{D}$ have a 3.8 fold increased risk of breast cancer compared to similar women in the highest quartile. The risk, OR (95% CI) was greater among women over the median age of 54, 5.2 (2.4, 11.1), than for younger women, 1.3 (0.7, 2.6).

Future work needs to address the temporal relationship between low levels of vitamin D metabolites and the development of breast cancer. A case-control study design cannot evaluate the temporal relationship; a prospective study would provide certainty on this question. A replication of this study using the preferred substrate of plasma or serum would allow a more accurate assessment of the $1,25(\text{OH})_2\text{D}$ levels which confer risk or protection.

Black women did not demonstrate the same relationship between low levels of $1,25(\text{OH})_2\text{D}$ and breast cancer risk. It is important to study a larger group of black women and other minority women to clarify the vitamin D/breast cancer relationship in this population.

The information on VDRs is important; differences in quantity and/or function may be associated with breast cancer risk or prognosis. Further, two recent papers described an association between Taq1 VDR genotype and prostate cancer risk^{13,14}.

The significance of this work relates to the potential for prevention of breast cancer. Recent work demonstrated the antineoplastic effect of a novel vitamin D analogue, 1α -hydroxyvitamin D_5 ¹⁵. Vitamin D analogues are potential chemotherapeutic as well as chemopreventive agents for breast cancer¹⁵. Finally, there is the possibility of better understanding the mechanisms of carcinogenesis as we define the role of vitamin D in this process.

References

1. DeLuca HF and Ostrem V. The relationship between the vitamin D system and cancer. *Adv Exp Med Biol* 206:413-429, 1987.
2. Reichel H, Koeffler HP, Norman AW. The role of the vitamin D endocrine system in health and disease. *NEJM* 320:980-991, 1989.
3. Manolagas SC. Vitamin D and its relevance to cancer. *AntiCancer Research* 7:625-638, 1987.
4. Garland CF, Garland FC, Shaw KO, Comstock GW, Helsing KJ, Gorham ED. Serum 25-hydroxyvitamin D and colon cancer: eight-year prospective study. *Lancet* 1176-1178, Nov. 18, 1989.
5. Corder EH, Guess HA, Hulka BS, Friedman GD, Sadler M, Vollmer RT, Lobaugh B, Drezner MK, Vogelmann JH, Orentreich N. Vitamin D and prostate cancer: a prediagnostic study with stored sera. *Cancer Epidemiology, Biomarkers & Prevention* 2:467-472, 1993.
6. Blot WJ, Fraumeni JF, Stone BJ. Geographic patterns of breast cancer in the United States. *J Natl Cancer Inst* 59:1407-1411, 1977.
7. Gorham ED, Garland FC, Garland CF. Sunlight and breast cancer incidence in the USSR. *Int J Epidemiol* 19:820-824, 1990.
8. Garland FC, Garland CF, Gorham ED, Young JF. Geographic variation in breast cancer mortality in the United States: a hypothesis involving exposure to solar radiation. *Preventive Med* 19:614-622, 1990.
9. Simard A, Vobecky J, Vobecky JS. Vitamin D deficiency and cancer of the breast: an unprovocative ecological hypothesis. *Can J Public Health* 82:300-303, 1991.
10. Jacobson EA, James K, Newmark HL, Carroll KK. Effects of dietary fat, calcium, and vitamin D on growth and mammary tumorigenesis induced by 7,12-dimethylbenz(a)anthracene in female Sprague-Dawley rats. *Cancer Res* 49:6300-6303, 1989.
11. Chouvet C, Vicard E, Devonec M, Saez S. 1,25-dihydroxyvitamin D₃ inhibitory effect on the growth of two human breast cancer cell lines (MCF-7, BT-20). *J Steroid Biochem* 24:373-376, 1986.
12. Frappart L, Falette N, Lefebvre MF, Bremond A, Vauzelle, JL, Saez S. In vitro studies of effects of 1,25-dihydroxyvitamin D₃ on the morphology of human breast cancer cell line BT-20. *Differentiation* 40:63-69, 1989.

13. Taylor JA, Hirvonen A, Watson M, Pittman G, Mohler JL, Bell DA. Association of prostate cancer with vitamin D receptor gene polymorphism. *Cancer Res*, 56:4108-4110, 1996.

14. Ingles SA, Ross RK, Yu MC, Irvine RA, La Pera G, Haile RW, Coetzee GA. Association of prostate cancer risk with genetic polymorphisms in vitamin D receptor and androgen receptor. *J Natl Cancer Inst*, 89:166-170, 1997.

15. Mehta RG, Moriarty RM, Mehta RR, Penmasta R, Lazzaro G, Constantinou A, Guo L. Prevention of preneoplastic mammary lesion development by a novel vitamin D analogue, 1α -hydroxyvitamin D₅.